



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

QUESTIONS AND DISCUSSIONS.

SEND ALL COMMUNICATIONS TO U. G. MITCHELL, University of Kansas, Lawrence.

NEW QUESTION.

34. Given the mixed integral and functional equation

$$\int_{x=0}^{x=x} f(x)dx = \frac{h}{6} \left[f(0) + 4f\left(\frac{x}{2}\right) + f(x) \right],$$

to determine the function $f(x)$. This equation is of rather fundamental practical value as it has to do with the most general solid whose volume is given by the prismatoid formula.

DISCUSSIONS.

I. CONCERNING AN ILLUSTRATION OF A CERTAIN NECESSARY CONDITION IN MINIMIZING A DEFINITE INTEGRAL WITH DISCONTINUOUS INTEGRAND.

By PAUL R. RIDER, Washington University.

The problem of minimizing a definite integral in which the integrand possesses a finite discontinuity along a plane curve has been considered by Bliss and Mason.¹ A certain necessary condition that they have discovered may be stated as follows:

If a curve C , of parameter t , which passes from the fixed point P_0 to the fixed point P_2 and crosses a curve D , whose equations are $x = x(\alpha)$, $y = y(\alpha)$, minimizes the sum of the two integrals

$$I = \int_{t_0}^{t_1} F(x, y, x', y')dt, \quad i = \int_{t_1}^{t_2} f(x, y, x', y')dt,$$

the first integral to be taken from the point P_0 to the curve D and the second from the curve D to the point P_2 , then at P_1 , the point of intersection of C and D , the relation

$$x_\alpha(F_{x'} - f_{x'}) + y_\alpha(F_{y'} - f_{y'}) = 0$$

must hold. The argument of x_α, y_α is the value of α for the point P_1 ; the arguments of $F_{x'}, F_{y'}$ are the values of x, y, x', y' on the curve C_{01} at the point P_1 , and those of $f_{x'}, f_{y'}$ are the values of the same variables on the curve C_{12} at the point P_1 .

This condition is well illustrated in a problem (Calculus 389) which recently appeared in the MONTHLY:

A man is at the southeast corner of a section of land and wishes to walk to the opposite corner in the least possible time. A circular track with a radius of $1/\pi$ miles is located in the section tangent to the west line at a point 120 rods from the south line. Conditions are such that he can walk at the rate of 4 miles an hour inside the track and 3 miles an hour outside the track. What course should he choose and how long is it? See the figure.

Solutions of the problem were given in the MONTHLY, Vol. 23, No. 24 (April, 1916), p. 125 by H. S. Uhler, and p. 127 by A. H. Holmes.

¹ BLISS and MASON, "A Problem of the Calculus of Variations in which the Integrand is Discontinuous," *Transactions of the American Mathematical Society*, Vol. 7 (1906), pp. 325-336.